

Datos Importantes

VIAGRA[®] (vi-AG-rah)
(sildenafil citrate) tablets

INFORMACIÓN IMPORTANTE SOBRE LA SEGURIDAD DE VIAGRA

Nunca tome VIAGRA si toma algún medicamento con nitratos. Esto incluye nitroglicerina. Su presión arterial podría disminuir rápidamente. Podría caer a un nivel inseguro o potencialmente mortal.

ACERCA DE LA DISFUNCIÓN ERÉCTIL (ED)

La disfunción eréctil significa que un hombre no puede tener o mantener una erección. La ED puede ser causada por problemas médicos, lesiones, o efectos secundarios de los fármacos. Es posible que no se conozca la causa.

ACERCA DE VIAGRA

VIAGRA es utilizado para tratar la ED en hombres. Cuando usted desee tener relaciones sexuales, VIAGRA puede ayudarle a tener y mantener una erección cuando está sexualmente excitado. Usted no puede tener una erección con sólo tomar la píldora. Sólo su médico puede indicarle VIAGRA. VIAGRA no cura la ED. VIAGRA no lo protege a usted ni a su pareja de las ETS (enfermedades de transmisión sexual) o el VIH. Deberá utilizar un condón. VIAGRA no es una hormona ni un afrodisíaco.

¿PARA QUIÉN ES VIAGRA?

¿Quién debe tomar VIAGRA?

Los hombres que sufran de ED y cuyo corazón tenga la salud suficiente para tener relaciones sexuales.

¿Quién NO debe tomar VIAGRA?

- Si alguna vez toma medicamentos con nitratos:
 - Medicamentos para tratar el dolor de pecho (angina), como la nitroglicerina o mononitrato o dinitrato de isosorbide
- Si consume algunas drogas de venta en la calle, como "poppers" (nitrato de amyl o nitrato)
- Si usted es alérgico a cualquiera de los ingredientes del comprimido de VIAGRA.

ANTES DE COMENZAR A TOMAR VIAGRA

Cuéntele a su médico si tiene o ha tenido:

- Ataque cardíaco, ritmo cardíaco anormal o infarto cerebral
- Problemas cardíacos, como insuficiencia cardíaca, dolores de pecho o estrechamiento de la válvula aórtica
- Presión arterial baja o alta
- Pérdida grave de la visión
- Una afección ocular llamada retinitis pigmentosa
- Tiene problemas renales o hepáticos
- Problemas sanguíneos, como anemia de células falciformes o leucemia
- Un pene deforme, enfermedad de Peyronie, o una erección que haya durado más de 4 horas
- Úlceras estomacales o cualquier tipo de problema hemorrágico

Cuéntele a su médico acerca de todos los medicamentos que tome.

- Incluya los medicamentos de venta sin receta médica, las vitaminas y los productos a base de hierbas. Cuéntele a su médico si toma o usa:
- Medicamentos llamados bloqueadores alfa para tratar la hipertensión o los problemas de próstata. Su presión arterial podría caer repentinamente demasiado bajo. Usted podría marearse o desmayarse. Su médico puede hacer que usted comience con una dosis más baja de VIAGRA.
 - Medicamentos llamados inhibidores de proteasa para el VIH. Su médico puede indicarle una dosis de 25 mg. Su médico puede limitar la dosis de VIAGRA a 25 mg en un periodo de 48 horas.
 - Otros métodos para causar erecciones. Estos incluyen píldoras, inyecciones, implantes o bombas.

POSIBLES EFECTOS SECUNDARIOS DE VIAGRA

Los efectos secundarios son principalmente leves a moderados. Estos efectos usualmente desaparecen después de unas pocas horas. Algunos de estos efectos tienen una mayor probabilidad de presentarse con las dosis más altas.

Los efectos secundarios más comunes son:

- Dolor de cabeza
- Sentirse sonrojado
- Malestar estomacal

Los efectos secundarios menos comunes incluyen:

- Problemas para distinguir entre el azul y el verde o ver tonos azules sobre las cosas
- Los ojos son más sensibles a la luz
- Visión borrosa

Rara vez, una cantidad reducida de hombres que tomaban VIAGRA informaron estos eventos adversos serios:

- Tener una erección que dura más de 4 horas. Si la erección no es tratada inmediatamente, podría presentarse una pérdida de potencia a largo plazo.
- Disminución repentina o pérdida de la agudeza visual en uno o ambos ojos. No sabemos si estos eventos son causados por el VIAGRA y los medicamentos como este, o son causados por otros factores. Dichos efectos secundarios pueden ser causados por afecciones como hipertensión o diabetes. Si usted presenta cambios repentinos en la visión, deje de utilizar VIAGRA y todos los medicamentos como este. Llame a su médico inmediatamente.
- Disminución repentina o pérdida de la audición. No sabemos si estos eventos son causados por el VIAGRA y los medicamentos como este, o son causados por otros factores. Si usted presenta cambios repentinos en la capacidad auditiva, deje de utilizar VIAGRA y todos los medicamentos como este. Llame a su médico inmediatamente.
- Ataque cardíaco, infarto cerebral, ritmo cardíaco irregular y muerte. No sabemos si estos eventos son causados por el VIAGRA o son causados por otros factores. La mayoría de estos efectos se presentaron en hombres que ya tenían problemas cardíacos.

Si presenta alguno de estos problemas, deje de tomar VIAGRA. Llame a su médico inmediatamente.

CÓMO TOMAR VIAGRA

Haga lo siguiente:

- Tome VIAGRA únicamente como su médico se lo indica. VIAGRA viene en comprimidos de 25 mg, 50 mg y 100 mg. Su médico le dirá cuál es la dosis adecuada para usted.
- Si tiene más de 65 años o tiene problemas hepáticos o renales graves, su médico puede comenzar con la dosis más baja (25 mg).
- Tome VIAGRA aproximadamente 1 hora antes de que desee tener relaciones sexuales. VIAGRA comienza a funcionar dentro de aproximadamente 30 minutos cuando está sexualmente excitado. El efecto de VIAGRA dura hasta 4 horas.

No haga lo siguiente:

- No tome VIAGRA más de una vez al día.
- No tome más VIAGRA del que su médico le indique. Si considera que necesita más VIAGRA, hable con su médico.
- No comience ni interrumpa ningún otro medicamento antes de consultar con su médico.

¿NECESITA MÁS INFORMACIÓN?

- Este es sólo un resumen de la información importante. Pídale a su médico o farmacéutico la información completa del producto O
- Visite www.viagra.com o llame al (888) 4-VIAGRA (484-2472).
- Los sitios web y los números telefónicos están disponibles actualmente sólo en inglés.
- Consulte la importante Información para el paciente al dorso del folleto.
- Consulte la Información de prescripción completa en www.pfizer.com.



¿Sin seguro? ¿Necesita ayuda para pagar las medicinas de Pfizer? Pfizer tiene programas que pueden ayudar. Llame al 1-866-706-2400 o visite www.PfizerHelpfulAnswers.com.



Distribuido por: Pfizer Labs
División de Pfizer Inc, NY, NY 10017
©2009 Pfizer Inc. Todos los derechos reservados.
Versión de marzo 2009 Impreso en los EE. UU.
Las marcas comerciales registradas son propiedad de sus respectivos dueños.
Sólo con receta médica VGU00488JP

PATIENT SUMMARY OF INFORMATION ABOUT **VIAGRA**[®]

(sildenafil citrate) tablets

This summary contains important information about VIAGRA[®]. It is not meant to take the place of your doctor's instructions. Read this information carefully before you start taking VIAGRA. Ask your doctor or pharmacist if you do not understand any of this information or if you want to know more about VIAGRA.

This medicine can help many men when it is used as prescribed by their doctors. However, VIAGRA is not for everyone. It is intended for use only by men who have a condition called erectile dysfunction. **VIAGRA must never be used by men who are taking medicines that contain nitrates of any kind, at any time. This includes nitroglycerin. If you take VIAGRA with any nitrate medicine your blood pressure could suddenly drop to an unsafe or life threatening level.**

• WHAT IS VIAGRA?

VIAGRA is a pill used to treat erectile dysfunction (impotence) in men. It can help many men who have erectile dysfunction get and keep an erection when they become sexually excited (stimulated).

You will not get an erection just by taking this medicine. VIAGRA helps a man with erectile dysfunction get an erection only when he is sexually excited.

• HOW SEX AFFECTS THE BODY

When a man is sexually excited, the penis rapidly fills with more blood than usual. The penis then expands and hardens. This is called an erection. After the man is done having sex, this extra blood flows out of the penis back into the body. The erection goes away. If an erection lasts for a long time (more than 6 hours), it can permanently damage your penis. You should call a doctor immediately if you ever have a prolonged erection that lasts more than 4 hours.

Some conditions and medicines interfere with this natural erection process. The penis cannot fill with enough blood. The man cannot have an erection. This is called erectile dysfunction if it becomes a frequent problem.

During sex, your heart works harder. Therefore sexual activity may not be advisable for people who have heart problems. Before you start any treatment for erectile dysfunction, ask your doctor if your heart is healthy enough to handle the extra strain of having sex. If you have chest pains, dizziness or nausea during sex, stop having sex and immediately tell your doctor you have had this problem.

• HOW VIAGRA WORKS

VIAGRA enables many men with erectile dysfunction to respond to sexual stimulation. When a man is sexually excited, VIAGRA helps the penis fill with enough blood to cause an erection. After sex is over, the erection goes away.

• VIAGRA IS NOT FOR EVERYONE

As noted above (*How Sex Affects the Body*), ask your doctor if your heart is healthy enough for sexual activity.

If you take any medicines that contain nitrates – either regularly or as needed – you should never take VIAGRA. If you take VIAGRA with any nitrate medicine or recreational drug containing nitrates, your blood pressure could suddenly drop to an unsafe level. You could get dizzy, faint, or even have a heart attack or stroke. Nitrates are found in many prescription medicines that are used to treat angina (chest pain due to heart disease) such as:

- nitroglycerin (sprays, ointments, skin patches or pastes, and tablets that are swallowed or dissolved in the mouth)
- isosorbide mononitrate and isosorbide dinitrate (tablets that are swallowed, chewed, or dissolved in the mouth)

Nitrates are also found in recreational drugs such as amyl nitrate or nitrite (“poppers”). If you are not sure if any of your medicines contain nitrates, or if you do not understand what nitrates are, ask your doctor or pharmacist.

VIAGRA is only for patients with erectile dysfunction. VIAGRA is not for newborns, children, or women. Do not let anyone else take your VIAGRA. VIAGRA must be used only under a doctor's supervision.

• WHAT VIAGRA DOES NOT DO

- VIAGRA does not cure erectile dysfunction. It is a treatment for erectile dysfunction.
- VIAGRA does not protect you or your partner from getting sexually transmitted diseases, including HIV—the virus that causes AIDS.
- VIAGRA is not a hormone or an aphrodisiac.

• WHAT TO TELL YOUR DOCTOR BEFORE YOU BEGIN VIAGRA

Only your doctor can decide if VIAGRA is right for you. VIAGRA can cause mild, temporary lowering of your blood pressure. You will need to have a thorough medical exam to diagnose your erectile dysfunction and to find out if you can safely take VIAGRA alone or with your other medicines. Your doctor should determine if your heart is healthy enough to handle the extra strain of having sex.

Be sure to tell your doctor if you:

- have ever had any heart problems (e.g., angina, chest pain, heart failure, irregular heart beats, heart attack or narrowing of the aortic valve)
- have ever had a stroke
- have low or high blood pressure
- have ever had severe vision loss
- have a rare inherited eye disease called retinitis pigmentosa
- have ever had any kidney problems
- have ever had any liver problems
- have ever had any blood problems, including sickle cell anemia or leukemia
- are allergic to sildenafil or any of the other ingredients of VIAGRA tablets
- have a deformed penis, Peyronie's disease, or ever had an erection that lasted more than 4 hours
- have stomach ulcers or any types of bleeding problems
- are taking any other medicines

• VIAGRA AND OTHER MEDICINES

Some medicines can change the way VIAGRA works. Tell your doctor about **any medicines** you are taking. Do not start or stop taking any medicines before checking with your doctor or pharmacist. This includes prescription and nonprescription medicines or remedies:

PATIENT SUMMARY OF INFORMATION ABOUT

VIAGRA®

(sildenafil citrate) tablets

- Remember, VIAGRA should never be used with medicines that contain nitrates (see *VIAGRA Is Not for Everyone*).
- If you are taking medicines called alpha-blockers for the treatment of high blood pressure or prostate problems, your blood pressure could suddenly drop. You could get dizzy or faint.
- If you are taking a protease inhibitor, your dose may be adjusted (please see *Finding the Right Dose for You*).
- VIAGRA should not be used with any other medical treatments that cause erections. These treatments include pills, medicines that are injected or inserted into the penis, implants or vacuum pumps.

• FINDING THE RIGHT DOSE FOR YOU

VIAGRA comes in different doses (25 mg, 50 mg and 100 mg). If you do not get the results you expect, talk with your doctor. You and your doctor can determine the dose that works best for you.

- Do not take more VIAGRA than your doctor prescribes.
- If you think you need a larger dose of VIAGRA, check with your doctor.
- VIAGRA should not be taken more than once a day.

Your doctor may prescribe a lower dose of VIAGRA in certain circumstances. For example:

- If you are older than age 65, or have serious liver or kidney problems, your doctor may start you at the lowest dose (25 mg) of VIAGRA.
- If you are taking protease inhibitors, such as for the treatment of HIV, your doctor may recommend a 25 mg dose and may limit you to a maximum single dose of 25 mg of VIAGRA in a 48 hour period.
- If you have prostate problems or high blood pressure for which you take medicines called alpha-blockers, your doctor may start you on a lower dose of VIAGRA.

• HOW TO TAKE VIAGRA

Take VIAGRA about 1 hour before you plan to have sex. Beginning in about 30 minutes and for up to 4 hours, VIAGRA can help you get an erection if you are sexually excited. If you take VIAGRA after a high-fat meal (such as a cheeseburger and french fries), the medicine may take a little longer to start working. VIAGRA can help you get an erection when you are sexually excited. You will not get an erection just by taking the pill.

• POSSIBLE SIDE EFFECTS

Like all medicines, VIAGRA can cause some side effects. These effects are usually mild to moderate and usually don't last longer than a few hours. Some of these side effects are more likely to occur with higher doses. The most common side effects of VIAGRA are headache, flushing of the face, and upset stomach. Less common side effects that may occur are temporary changes in color vision (such as trouble telling the difference between blue and green objects or having a blue color tinge to them), eyes being more sensitive to light, or blurred vision.

PATIENT SUMMARY OF INFORMATION ABOUT

VIAGRA®

(sildenafil citrate) tablets

In rare instances, men taking PDE5 inhibitors (oral erectile dysfunction medicines, including VIAGRA) reported a sudden decrease or loss of vision in one or both eyes. It is not possible to determine whether these events are related directly to these medicines, to other factors such as high blood pressure or diabetes, or to a combination of these. If you experience sudden decrease or loss of vision, stop taking PDE5 inhibitors, including VIAGRA, and call a doctor right away.

In rare instances, men have reported an erection that lasts many hours. You should call a doctor immediately if you ever have an erection that lasts more than 4 hours. If not treated right away, permanent damage to your penis could occur (see *How Sex Affects the Body*).

Sudden loss or decrease in hearing, sometimes with ringing in the ears and dizziness, has been rarely reported in people taking PDE5 inhibitors, including VIAGRA. It is not possible to determine whether these events are related directly to the PDE5 inhibitors, to other diseases or medications, to other factors, or to a combination of factors. If you experience these symptoms, stop taking VIAGRA and contact a doctor right away.

Heart attack, stroke, irregular heart beats, and death have been reported rarely in men taking VIAGRA. Most, but not all, of these men had heart problems before taking this medicine. It is not possible to determine whether these events were directly related to VIAGRA.

VIAGRA may cause other side effects besides those listed on this sheet. If you want more information or develop any side effects or symptoms you are concerned about, call your doctor.

• ACCIDENTAL OVERDOSE

In case of accidental overdose, call your doctor right away.

• STORING VIAGRA

Keep VIAGRA out of the reach of children. Keep VIAGRA in its original container. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

• FOR MORE INFORMATION ON VIAGRA

VIAGRA is a prescription medicine used to treat erectile dysfunction. Only your doctor can decide if it is right for you. This sheet is only a summary. If you have any questions or want more information about VIAGRA, talk with your doctor or pharmacist, visit www.viagra.com, or call 1-888-4VIAGRA.



LAB-0220-6.0

October 2007